

DONJOY®

Playmaker II FourcePoint

I Can
Play

Megan Rapinoe
U.S. Soccer Player
Olympic Gold Medalist

 **DJO**
GLOBAL



Women are **2^{TO}10X** more likely to injure their ACL¹

A person who has *torn their ACL* has



15X greater risk of a second ACL injury²

(during initial 12-months after ACL reconstruction)



50% of ACL injuries occur in **15-25** year olds³

60%^{TO} 80% of ACL injuries are **non-contact** related⁴

Risk of ACL injury to the **opposing knee**
(opposite knee) is

2X



that of the **reconstructed knee**⁵



Psychological factors, primarily

fear of re-injury,
influence the ability to return
to play⁶

I Can Recover

Post-injury with
FourcePoint Hinge Technology

Increases flexion angles
(minimizes time spent
in "at risk" position)⁷




Improves joint mechanics
in BOTH surgical and
non-surgical knees⁸



Immeasurable Comfort
Hino™ (high tenacity nylon/
spandex fabric - posterior) offers
unmatched comfort and fit.




Just Plain "Easy"
Molded strap tabs allow quick
and easy strap adjustments
to deliver a secure and
comfortable fit.



"Cooler" Experience
Advanced spacer fabric
promotes a "cool" environment
for activities on all levels.





I Can Train Both Knees with One Brace

Preventative use for non-contact sports with
FourcePoint Hinge Technology

60% decrease in
non-contact ACL injury rate in
BOTH knees (one brace protects both knees)⁹

Training effects
retained by about
50% while
not wearing
the brace⁹





ORDERING INFORMATION	
11-3504-1	PLAYMAKER II FP, XS
11-3504-2	PLAYMAKER II FP, S
11-3504-3	PLAYMAKER II FP, M
11-3504-4	PLAYMAKER II FP, L
11-3504-5	PLAYMAKER II FP, XL
11-3504-6	PLAYMAKER II FP, XXL
11-3504-7	PLAYMAKER II FP, XXXL

Call 800.336.6569 or visit www.djoglobal.com/donjoy  

“I have had two major injuries during my career—both ACL tears. There is a lot of pressure inside and outside of schools and organizations to recover quickly, which I appreciate, but it’s up to young athletes, parents and coaches to take steps to recover safely and guard against injury and re-injury. This will only allow for a stronger, better return to sport in the long run.”

- Megan Rapinoe, U.S. Soccer Player and Olympic Gold Medalist

1. Silvers HJ, Mandelbaum BR. Prevention of Anterior Cruciate Ligament Injury in the Female Athlete. *Brj SportsMed*. 2007; 41(Suppl): 152-159.

2. Paterno MV. Paper #2. Presented AOSM Annual Meeting; July 11-14, 2013; Chicago.

3. Griffin LY, Albohm MJ, Arendt, EA, et al. Understanding and Preventing Noncontact Anterior Cruciate Ligament Injuries: A Review of the Hunt Valley II Meeting, January 2005. *American Journal of Sports Medicine*. 2006; 34:9. 1512-1532.

4. Arendt EA, Agel J, Dick R. Anterior cruciate ligament injury patterns among collegiate men and women. *Journal of Athletic Training*, 1999; 34(2):86-92.

5. Wright RW, Magnussen RA, Dunn WR, Spindler KP. Ipsilateral Graft and Contralateral ACL Rupture at Five Years or More Following ACL Reconstruction. *Journal of Bone and Joint Surgery, Am*. 2011;93:1159-1165.

6. Spindler et al. Return to High School and College-Level Football After Anterior Cruciate Ligament Reconstruction. *Am J Sports Med*. Published online Aug. 24, 2012.

7. Yu B, Herman D, Preston J, Lu W, Kirkendall DT, Garrett WE. Immediate Effects of a Knee Brace with a Constraint to Knee Extension on Knee Kinematics and Ground Reaction Forces in a Stop-Jump Task. *AJSM*. 2004; 32:1136-1143.

8. Queen R, Butler RJ, Dai B, Garrett WE. Effects of Knee Extension Constraint Bracing on Lower Extremity Motion Patterns in Post-ACL Reconstruction Patients. Interim Report of 6-12 month data from the ongoing study, 2012

9. Yu B. Training Effects of a Knee Extension Constraint Brace on Lower Extremity Motion Patterns and the Risk for Non-Contact ACL Injury. Feagin Leadership Forum. Duke Sports Medicine Center, 2010.



DJO, LLC | A DJO Global Company

T 800.336.6569 D 760.727.1280 F 800.936.6569

1430 Decision Street | Vista, CA 92081-8553 | U.S.A.

DJOGlobal.com/donjoy

Together in Motion™

DJO Global proudly sponsors:

